arXiv:1205.4597v1 [math.AT] 21 May 2012

COMPUTATIONS IN RATIONAL SECTIONAL CATEGORY

J.G. CARRASQUEL VERA

Abstract. We give a simple upper bound for the rational sectional category and use it to compute invariants of the type of Farber's topological complexity of rational spaces. We also show that the sectional category of formal morphisms reaches its cohomological lower bound and that the higher topological complexity of a formal space is determined by its Lusternik-Schnirelmann category and its topological complexity.

1. Introduction

This paper concerns the rational sectional category of a continuous map $f : X \to Y$ and in particular the rational topological complexity of a space X. All the spaces considered will be supposed simply connected with finite Betti numbers.

Recall that the sectional category of f, $secat(f)$, is the least integer k for which there is an open cover $\{U_0, \dots U_k\}$ of Y and maps $s_i : U_i \to X$ such that $f \circ s_i$ is homotopic to the inclusion of U_i in Y. When X is a point, the sectional category of f is the usual Lusternik-Schnirelmann category of Y. [1]

Denote by $H^*(X; R)$ the cohomology ring of X with coefficients in a ring R. It is well known ([1]) that

$$secat(f) \geq nil\ Ker(p^*),$$

where *nil* denotes the *nilpotence* of an ideal,

$$nil\ I = \max\{k :\ \exists x_1 \dots x_k \in I \text{ with } x_1 \cdots x_k \neq 0\}.$$

Denote X_0 the rationalization of X and $f_0 : X_0 \to Y_0$ the rationalization of f, our main result is an upper bound for $secat(f_0)$. Let $(A, d) \to (B, d)$ be a surjective model of f_0 ([4]) with kernel K. Then, we prove the following which generalizes [6, Prop. 1.3],

Theorem 1.1. *Let m be the minimum k such that the quotient map*

$$(A, d) \to \left(\frac{A}{K^{k+1}}, \overline{d}\right)$$

admits a homotopy retraction, then $m \geq secat(f_0)$.

We can use this result for computations of the higher topological complexity of a space X, $TC_n(X)$. For recall $TC_n(X) = secat(\Delta_n)$ where $\Delta_n : X \to X^n$ is the $n-$diagonal map. In [2], Farber proves that

$$cat(X) \leq TC(X) \leq 2cat(X).$$

1991 *Mathematics Subject Classification.* Primary 55P62, Secondary 55M30.
Key words and phrases. Topological complexity, Rational homotopy, Sectional category.
This work was partially supported by the Spanish grant MTM2010-18089.

Corollary 1.2. *If X is a formal space, then*

$$TC_n(X_0) = nil\ Ker\left(\Delta^* : H^*(X^n;\mathbb{Q}) \to H^*(X;\mathbb{Q})\right).$$

Corollary 1.3. *If $\pi_q(X)$ is a finite group for all q except for a finite number of odd integers $q_1, \ldots, q_r$, then $TC_n(X_0) = (n-1)cat(X_0)$.*

Corollary 1.4. *Let X be a formal space, then for $n \geq 3$*

$$TC_n(X_0) = TC(X_0) + (n-2)cat(X_0).$$

2. Sectional category, topological complexity and rational homotopy.

Sectional category was first defined by A.S. Schwarz in [9]. A good exposition can be found in [1], section 9.3.

Definition 2.1. Let $f : X \to Y$ be a map. The *sectional category* of f, $secat(f)$, is the least integer k for which there is an open cover $\{U_0, \ldots, U_k\}$ of Y and maps $s_i : U_i \to X$, such that $p \circ s_i$ is homotopic to the inclusion of U_i in Y.

The *topological complexity* of X is the sectional category of the map $p : X^{[0,1]} \to X \times X$, $p(\omega) = (\omega(0), \omega(1))$. This is the same as the sectional category of the diagonal map of X. This notion has been introduced by M. Farber in [2] and explicit computations can be found in [6] and [7], for instance.

Denote X_0 the rationalization of X and $f_0 : X_0 \to Y_0$ the rationalization of f. Then ([1]) $secat(f_0) \leq secat(f)$, and in particular $TC(X_0) \leq TC(X)$.

Every simply connected space X with finite Betti numbers has a (Sullivan) minimal model $(\Lambda V, d)$. This is a commutative differential graded algebra (**CDGA** for short), where ΛV denotes the free graded commutative algebra on a graded vector space V and where $d(V) \subset \Lambda^{\geq 2}V$, see [4]. Moreover, each morphism of **CDGA** $\varphi : (A, d) \to (B, d)$ admits a minimal relative model



where i is the canonical injection, $\psi \circ i = \varphi$,

$$d(V) \subset \left(A^+ \otimes \Lambda V\right) + \left(A \otimes \Lambda^{\geq 2}V\right),$$

and ψ is a quasi-isomorphism.

Now we transform i into a surjective map

$$h : \left(A \otimes \Lambda V \otimes \Lambda \hat{V}, D\right) \to (A \otimes \Lambda V, d)$$

in a *standard way* by putting

$\hat{V}^n = V^{n-1}$, $D(\hat{V}) = 0$, $D(v) = \hat{v}$, $h(a) = a$, $h(v) = v$ and $h(\hat{v}) = d(v)$.

Suppose $f : X \to Y$ is a cofibration. Denote by $T^{m+1}(f)$ the fat wedge of f,

$$T^{m+1}(f) = \{(y_0, \ldots, y_m) : \ y_i \in f(X) \text{ for at least one } i\} \subset Y^{m+1},$$

and by $\beta : (\Lambda V, d)^{\otimes m+1} \to (\Lambda Z, d)$ a model for the injection $T^{m+1}(f) \hookrightarrow Y^{m+1}$.

Definition 2.2. $Secat_0(f)$ is the smallest m for which there exists a **CDGA** morphism $\rho : (\Lambda Z, d) \to (\Lambda V, d)$ such that $\rho \circ \beta \sim \mu_{m+1}$,

$$\begin{array}{ccc} (\Lambda V, d)^{\otimes m+1} & \xrightarrow{\mu_{m+1}} & (\Lambda V, d) \\ {\scriptstyle\beta}\downarrow & \nearrow_{\rho} & \\ (\Lambda Z, d) & & \end{array}$$

Where μ_{m+1} denotes the multiplication map.

Then,

Theorem 2.3 ([3], Chap. 8)**.** $Secat_0(f) = Secat(f_0)$.

From this Theorem, Fassò-Velenik in [3] deduces the following more explicit characterisation

Theorem 2.4 ([3], chap. 8)**.** *Let $\overline{f} : (\Lambda V, d) \to (\Lambda V \otimes \Lambda W, d)$ be a relative Sullivan model for f and $h : \left(\Lambda V \otimes \Lambda W \otimes \Lambda \hat{W}, D\right) \to (\Lambda V \otimes \Lambda W, d)$ be the standard surjective extension of $\overline{f}$. The $secat(f_0)$ is the smallest m such that the morphism*

$$(\Lambda V, d) \longrightarrow \frac{\bigotimes_{\Lambda V}^{m+1} \left(\Lambda V \otimes \Lambda W \otimes \Lambda \hat{W}\right)}{\bigotimes_{\Lambda V}^{m+1} Ker\ h}$$

admits a homotopy retraction.

3. Homotopy retractions and another characterization of $secat(f_0)$

Definition 3.1. Let $f : A \to B$ be a **CDGA** morphism. A *homotopy retraction* for f is a morphism $r : (A \otimes \Lambda V, D) \to A$ such that $r \circ i = Id_A$ where $i : A \hookrightarrow (A \otimes \Lambda V, D)$ is a relative Sullivan model for f.

Lemma 3.2. *Consider the following commutative* **CDGA** *diagram.*

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ {\scriptstyle\varphi}\downarrow{\scriptstyle\simeq} & & {\scriptstyle\simeq}\downarrow{\scriptstyle\psi} \\ C & \xrightarrow[g]{} & D \end{array}$$

Then there exists a homotopy retraction for f if and only if there exists one for g.

Proof. Take $(A \otimes \Lambda V, D)$ a relative Sullivan model for f and define $(C \otimes \Lambda V, D) = C \otimes_A (A \otimes \Lambda V, D)$. The diagram



tells us that $C \otimes \Lambda V$ is a relative Sullivan model for g (see [4, Pg. 183] for details). Observe that the quasi-isomorphism $C \otimes \Lambda V \xrightarrow{\simeq} D$ is given by pushout's universal property.

Suppose there exists a homotopy retraction r_1 for f. Then

$$r_2 = Id \otimes_A r_1 : C \otimes_A (A \otimes \Lambda V) \to C \otimes_A A$$

is a homotopy retraction for g. Now if r_2 exists, the relative lifting lemma ([4, Pg. 186]) gives birth to r_1 as follows:



□

Lemma 3.3. *Consider the following commutative* **CDGA** *diagram.*



If β admits homotopy retraction, so does α.

Proof. Take relative Sullivan models for α and β. We have the following commutative diagram



The lifting lemma ([4, Pg. 186]) gives a map h



making commutative the diagram



Write $r : (B \otimes \Lambda W, D) \to B$ a homotopy retraction for β. The desired homotopy retraction $s : (A \otimes \Lambda V) \to A$ for α is given by the lifting lemma in



□

Now let $f : A \to B$ be a **CDGA** morphism. We consider the diagram



where

(i) $\theta : (A \otimes \Lambda V, D) \xrightarrow{\simeq} B$ is a minimal relative Sullivan model for f and h its standard surjective replacement,
(ii) i and j are standard inclusions.

Lemma 3.4. *Every differential ideal M of $\left(A \otimes \Lambda V \otimes \Lambda \hat{V}, D\right)$, that is free as A-module, $M = A \otimes W$, is a semifree A-module.*

Proof. For each r, define

$$W(r) := W \cap \bigoplus_{q+2p \leq r} \left(\Lambda^p V \otimes \Lambda \hat{V}\right)^q.$$

Since $D\left(\left(\Lambda^p V \otimes \Lambda \hat{V}\right)^q\right) \subset \left(\Lambda^{p-1} V \otimes \Lambda \hat{V}\right)^{q+1}$, $D(W(r)) \subset W(r-1)$ and, thus $\{A \otimes W(r)\}$ is a semifree filtration for $A \otimes W$. □

Proposition 3.5. *With previous notation, $secat(f_0)$ is the smallest m for which the map*

$$p_{m+1} : A \longrightarrow \frac{\otimes_A^{m+1}\left(A \otimes \Lambda V \otimes \Lambda \hat{V}\right)}{\otimes_A^{m+1} Ker(h)}$$

admits a homotopy retraction.

Proof. Let $k : (\Lambda Z, d) \xrightarrow{\simeq} (A, d)$ be a minimal model for (A, d) and $(\Lambda Z \otimes \Lambda W, D)$ be a minimal relative model for $f \circ k : (\Lambda Z, d) \to (B, d)$.

Then $(A \otimes \Lambda W, D) := (A, d) \otimes_{\Lambda Z} (\Lambda Z \otimes \Lambda W, D)$ is a minimal relative model for $f : (A, d) \to (B, d)$. Since the minimal relative model is unique up to isomorphism, we can suppose $V = W$.

Consider the following diagram where $(A \otimes \Lambda V, D) := A \otimes_{\Lambda Z} (\Lambda Z \otimes \Lambda V, D)$ and where the top and bottom triangles are the standard surjective decompositions for j' and j,



By Lemma 3.4, $Ker(h)$ is A-semifree and $Ker(h')$ is ΛZ-semifree. The five lemma guarantees a quasi-isomorphism $Ker(h') \xrightarrow{\simeq} Ker(h)$ which induces (induction and [4, Th. 6.7 ii)]) a quasi-isomorphism $\otimes^m_{\Lambda Z} Ker(h') \xrightarrow{\simeq} \otimes^m_A Ker(h)$. Again, the five lemma gives rise to quasi-isomorphisms

$$\begin{array}{ccc} \Lambda Z & \xrightarrow{p'_m} & \frac{\otimes^m_{\Lambda Z}\left(\Lambda Z \otimes \Lambda V \otimes \Lambda \hat{V}\right)}{\otimes^m_{\Lambda Z} Ker(h')} \\ \simeq \downarrow & & \downarrow \simeq \\ A & \xrightarrow[p_m]{} & \frac{\otimes^m_A\left(A \otimes \Lambda V \otimes \Lambda \hat{V}\right)}{\otimes^m_A Ker(h)}. \end{array}$$

Lemma 3.2 tells us that p'_m admits homotopy retraction if and only if p_m admits homotopy retraction, but Theorem 2.4 tells us that $secat_0(f)$ is the least m for which p'_{m+1} admits a homotopy retraction. □

4. Upper bound for $secat_0$

Lemma 4.1. *Let $f : A \to B$ be a surjective* **CDGA** *morphism. For each $[a] \in \ker(H(f))$ there exists $a' \in A$ such that $a - da' \in \ker(f)$.*

Proof. Let $[a] \in \ker(H(f))$, this means that $da = 0$ and $f(a) = db$ for certain $b \in B$. Since f is surjective, there is $a' \in A$ such that $f(a') = b$. So $0 = f(a) - db = f(a) - df(a') = f(a - da')$ and thus $a - da' \in \ker(f)$.

□

Lemma 4.2. *Let $f : A \to B$ be a surjective* **CDGA** *morphism such that $H(f)$ is also surjective. Then there is a minimal relative model for f, $\theta : (A \otimes \Lambda V, D) \xrightarrow{\simeq} B$ such that*

$$D(V) \subset (K \otimes 1) \oplus \left(A \otimes \Lambda^+ V\right),$$

where K denotes the kernel of f.

Proof. Since $H(f)$ is surjective, we define $V^{\leq k}$ and

$$\theta_k : \left(A \otimes \Lambda V^{\leq k}, D\right) \longrightarrow (B, d)$$

by induction on k with the following properties

(i) $H^{\leq k+1}(\theta_k)$ is injective,
(ii) $D(V^{\leq k}) \subset (K \otimes 1) \oplus \big(A \otimes \Lambda^+ V^{<k}\big)$,
(iii) $\theta_k(V^{\leq k}) = 0$.

Suppose we have constructed θ_k, we will construct θ_{k+1} inductively. Write $V^{k+1} = \bigoplus_{r\geq 0} W_r$ and define $\theta_{k,r} : \big(A \otimes \Lambda V^{\leq k} \otimes W_{\leq r}, D\big) \longrightarrow (B, d)$ with the following properties

(a) If $[u] \in Ker\ H^{k+2}(\theta_{k,r-1})$ then $[u] = 0$ in $\big(A \otimes \Lambda V^{\leq k} \otimes W_{\leq r}, D\big)$,
(b) $D(V^{\leq k} \oplus W_{\leq r}) \subset (K \otimes 1) \oplus \big(A \otimes \Lambda^+(V^{\leq k} \oplus W_{<r})\big)$,
(c) $\theta_{k,r}(V^{\leq k} \oplus W_{\leq r}) = 0$.

Suppose $\theta_{k,r}$ has been defined. From (c) we deduce that $Ker\ \theta_{k,r} = (K \otimes 1) \oplus \big(A \otimes \Lambda^+(V^{\leq k} \oplus W_{\leq r})\big)$. Now let $\{[\xi_\alpha]\}_{\alpha\in\Gamma}$ be a basis for $Ker\ H^{k+2}(\theta_{k,r})$. By the previous lemma, there exists $\varphi_\alpha \in (A \otimes \Lambda V_{\leq k} \otimes \Lambda W_{\leq r})$ such that $\xi_\alpha - D(\varphi_\alpha) \in Ker\ \theta_{k,r}$. Define W_{r+1} with basis $\{w_\alpha\}_{\alpha\in\Gamma}$ and extend $\theta_{k,r}$ and D by $\theta_{k,r+1}(w_\alpha) = 0$ and $D(w_\alpha) = \xi_\alpha - D(\varphi_\alpha)$. Note that the minimality of the model is assured by degree reasons since $deg(\xi_\alpha) = k + 2$ and $V^{\leq k} \oplus W_{\leq r}$ is concentrated in degrees smaller than or equal to $k + 1$. □

Now let $f : A \to B$ be any surjective **CDGA** morphism with kernel K. For each class $[u_\alpha] \in coker\ H(f)$, we choose an element $a_\alpha \in A$ with $f(a_\alpha) = u_\alpha$ and we introduce a new generator v_α with $deg(v_\alpha) = deg(u_\alpha)$. We put $Z = \{v_\alpha\}$ and extend f to

$$g : (A \otimes \Lambda Z, D) \longrightarrow (B, d)$$

by $g(v_\alpha) = u_\alpha$ and $D(v_\alpha) = 0$.

Then g is a surjective **CDGA** morphism such that $H(g)$ is also surjective. We apply Lemma 4.2 and deduce:

Lemma 4.3. *Let $f : A \to B$ be a surjective* **CDGA** *morphism. Then with the notation above, f admits a minimal relative Sullivan model*

$$\theta : (A \otimes \Lambda Z \otimes \Lambda W, D) \longrightarrow (B, d)$$

such that $D(Z \oplus W)$ is in the ideal generated by K, W and the set $\{v_\alpha - a_\alpha\}$.

We can now prove our main theorem.

Theorem 4.4. *Let $f : A \twoheadrightarrow B$ be a surjective* **CDGA** *morphism with K denoting its kernel. If the projection*

$$\rho_{m+1} : A \to A/K^{m+1}$$

admits an homotopy retraction then $secat_0(f) \leq m$.

Proof. Fix m. We will prove that if ρ_m admits homotopy retraction then also does the map

$$p_m : A \longrightarrow \frac{\otimes_A^m \left(A \otimes \Lambda V \otimes \Lambda \hat{V}\right)}{\otimes_A^m Ker(h)}.$$

Denote by $\pi : A \otimes \left(\Lambda V \otimes \Lambda \hat{V}\right)^{\otimes m} \to A$ be the projection defined as follows. Write $V = Z \otimes W$ as in Lemma 4.3. Then put $\pi(w) = \pi(\hat{w}) = 0$, and for $v_\alpha \in Z$,

$\pi(v_\alpha) = a_\alpha$, $\pi(\hat{v}_\alpha) = d(a_\alpha) \in K$. Observe that π is a quasi-isomorphism.

Since $Ker\ h$ is generated by the $\hat{v}_\alpha$ and the $\hat{w} - D(w)$, $\pi(Ker\ h) \subset K$. We obtain the following commutative diagram where $\overline{\pi}$ is well defined.



If ρ_m admits homotopy retraction, Lemmas 3.2 and 3.3 tell us that also p_m does. □

5. Applications

Definition 5.1. Let $f : A \to B$ a **CDGA** morphism. We say that f is *formal* when there is a commutative **CDGA** diagram



A continuous map is said to be *formal* if it admits a formal **CDGA** model.

For more on formal morphisms the reader is referred to [5] and [11]. It is obvious from this definition that if $f : A \to B$ is formal then both A and B are formal as well.

Example 5.2. Let A be a formal **CDGA** , clearly the m-multiplication morphism $A^{\otimes m} \to A$ is formal morphism. Therefore, if X is formal, then the n-diagonal map $\Delta_X^n : X \hookrightarrow X^n$ is also formal.

Theorem 5.3. *Let $f : A \to B$ be a formal morphism with $H(f)$ surjective. Then*

$$secat(f) = nil(kerH(f)).$$

Proof. By formality, $secat(f) = secat(H(f))$. Write $m := nil(kerH(f))$, we must prove that $secat(H(f)) \leq m$ but this is direct consequence of Theorem 4.4 and the fact that $(KerH(f))^{m+1} = \{0\}$. □

The following definition was given by Y.B Rudyak in [8].

Definition 5.4. Let X be a topological space and $\Delta_X^n : X \hookrightarrow X^n$ its n diagonal map. The *n-topological complexity* of X is the sectional category of the map Δ_X^n, $TC_n(X) = secat(\Delta_X^n)$.

Observe that the case $n = 2$ yields the well known topological complexity introduced by M. Farber in [2].

Proposition 5.5. *Let X be a formal space and denote $\mu_n : H^*(X,\mathbb{Q})^{\otimes n} \twoheadrightarrow H^*(X,\mathbb{Q})$ the n-multiplication of $H^*(X,\mathbb{Q})$. Then $TC_n(X_0) = nil\,(Ker(\mu_n))$.*

Proof. If X is a formal space, Δ_X^n is a formal map modelled by μ_n. Since rationalization commutes with limits, we have that $TC_n(X_0) = secat(\Delta^n_{X_0}) = secat((\Delta^n_X)_0)$. The result follows from Theorem 5.3. □

Observe that in the case $n = 2$, last proposition is just the main result in [7].

We will often use the following two lemmas.

Lemma 5.6. *Let X be a subset of the* **CDGA** *A and I the ideal of A generated by X. Then nil I is the largest m for which there exist $x_1,\dots,x_m \in X$ such that $x_1\cdots x_m \neq 0$.*

Lemma 5.7. *Let A be a* **CDGA** *such that A^+ is generated by $\{x_i\}_{i\in I}$. Then K_n, the kernel of the n-multiplication morphisms $\mu_n : A^{\otimes n} \to A$, is generated by $\{x_{i,j} - x_{i,1} :\ i \in I,\ 2 \le j \le n\}$ where*

$$x_{i,j} = 1\otimes\cdots\otimes 1\otimes x_i\otimes 1\otimes\cdots\otimes 1 \in A^{\otimes j-1}\otimes A^+\otimes A^{\otimes n-j}.$$

Theorem 5.8. *Let A be a* **CDGA** *and K_n the kernel of the n-multiplication morphism $A^{\otimes n}\to A$. Then for $n \ge 3$,*

$$nil\ K_n = nil\ K_{n-1} + nil\ A^+.$$

Proof. Write $r = nil\ K_{n-1}$ and $s = nil\ A^+$. Consider $\omega \neq 0$ a product of r factors in K_{n-1} and $\alpha = a_1\cdots a_s \neq 0$ with $a_i \in A^+$. Then the element $(\omega\otimes 1)(a_{1,n} - a_{1,1})\cdots(a_{s,n} - a_{s,1}) = (\omega\otimes\alpha) + \xi$ with $\xi \in A^{\otimes n-1}\otimes A^{<deg(\alpha)}$ must be non-zero. This proves that $nil\ K_n \ge r + s$.

On the other hand, write $m = nil\ K_n$. Then there exists $\omega \neq 0$ a product of m factors in K_n which can be written as

$$\omega = (P\otimes 1)\cdot(x_{i_1,n} - x_{i_1,1})^{n_1}\cdots(x_{i_t,n} - x_{i_t,1})^{n_t},$$

with P a product of factors in K_{n-1}, $0 \le n_i$ and the x_{i_j} pairwise different. Then

$$\omega = \sum_{\sigma\in\Sigma}\alpha_\sigma\otimes\left(\prod_{j=1}^{t} x_{i_j,n}^{\sigma_j}\right)$$

with $\Sigma = \{(\sigma_1,\dots,\sigma_t) :\ 0 \le \sigma_j \le n_j\}$. Chose $\sigma\in\Sigma$ such that $\alpha_\sigma\otimes\left(\prod_{j=1}^t x_{i_j,n}^{\sigma_j}\right)$ is non-zero and $\sum_j \sigma_j$ maximal. In this case, $\sum_j\sigma_j \le nil\ A^+$ and

$$\alpha_\sigma = \pm P\cdot\left(\prod_{j=1}^{t}\binom{n_j}{\sigma_j}x_{i_j,1}^{n_j-\sigma_j}\right) \neq 0.$$

Now decompose $\alpha_\sigma \in A\otimes A^{\otimes n-2}$ as

$$\alpha_\sigma = \sum_{u\in U} w_u\otimes w'_u + \rho,$$

with $\sum_{u\in U} w_u \otimes w'_u \neq 0$, $deg(w_u) = l$, $w'_u \in A^{\otimes n-2}$, and $\rho \in A^{<l} \otimes A^{\otimes n-2}$. Consider the element

$$\beta = P \cdot \prod_{j=1}^{t} \binom{n_j}{\sigma_j} \left(x_{i_j,2} - x_{i_j,1}\right)^{n_j-\sigma_j} = \sum_{u\in U} w_u \otimes w'_u + \rho',$$

with $\rho' \in A^{<l} \otimes A^{\otimes n-2}$. Note that β is a non-zero product of $m - \sum_j \sigma_j$ factors in K_{n-1}. This proves that $nil\ K_n = m \leq nil\ K_{n-1} + \sum_j \sigma_j \leq nil\ K_{n-1} + nil\ A^+$. □

Corollary 5.9. *Let X be a formal space, then for $n \geq 3$*

$$TC_n(X_0) = TC_{n-1}(X_0) + cat(X_0) = TC(X_0) + (n-2)cat(X_0).$$

Example 5.10. Let X be the wedge $(S^3 \times S^3) \vee (S^3 \times S^3)$. Then $TC(S^3 \times S^3) = 2$ but $TC(X) = 4$.

The cohomology of X is $\Lambda(x, y, z, t)/(xz, xt, yz, yt)$, with x, y, z, t in degree 3. Then writing $x^- = 1\otimes x - x\otimes 1$ and so on, we see that $x^-y^-z^-t^- = xy\otimes zt + zt\otimes xy$. This shows that $nil\ Ker\ \mu_2 = 4$.

More generally, let X and Y be formal spaces, then $TC(X \vee Y) \geq cat(X) + cat(Y)$.

Indeed, let w_1 and w_2 be monomials in $H^*(X)$ and $H^*(Y)$ of maximal length n and m, $w_1 = a_1 \cdots a_n$ and $w_2 = b_1 \cdots b_m$. Then the identity

$$a_1^- \cdots a_n^- b_1^- \cdots b_m^- = w_1 \otimes w_2 \pm w_2 \otimes w_1$$

shows that $TC(X \vee Y) \geq n + m$.

Theorem 5.11 ([6], Thm. 1.4)**.** *Suppose $\pi_*(X) \otimes \mathbb{Q}$ is finite dimensional and concentrated in odd degrees. Then $TC(X_0) = cat(X_0)$.*

Proof. By hypothesis a model of X has the form $(A, d) = (\Lambda(x_1, \ldots, x_r), d)$ with x_i in odd degree. Then the kernel K of the multiplication $A \otimes A \to A$ is generated by the elements $x_i \otimes 1 - 1 \otimes x_i$. Since the square of these elements is zero, $K^{r+1} = 0$. This implies that $cat(X_0) \leq TC(X_0) \leq r = cat(X_0)$. □

It follows more generally,

Theorem 5.12. *If $\pi_*(X) \otimes \mathbb{Q}$ is finite dimensional and concentrated in odd degrees. Then $TC_n(X_0) = (n-1)cat(X_0)$.*

Proof. Let $(A, d) = (\Lambda(x_1, \ldots, x_r), d)$ be a model of X with x_i in odd degree. Then

$$(A, d)^{\otimes n} = (\Lambda(x_{1,1}, \ldots, x_{r,1}, x_{1,2}, \ldots, x_{2,r}, \ldots, x_{1,n}, \ldots, x_{r,n}), d)$$

and K the kernel of the n multiplication is generated by the elements

$$\{x_{i,j} - x_{i,1} : \ 1 \leq i \leq r, \ 2 \leq j \leq n\}.$$

Since the square of these elements is zero, we have that $K^{r(n-1)+1} = 0$.

Now consider the following pullback diagram

$$\begin{array}{ccc} PX & \longrightarrow & X^{[0,1]} \\ {\scriptstyle q}\downarrow & & \downarrow{\scriptstyle p} \\ X^{n-1}\times * & \longrightarrow & X^n \end{array}$$

where $p(\omega) = \Big(\omega(0), \omega(\frac{1}{n-1}), \omega(\frac{2}{n-1}), \dots, \omega(1)\Big)$. Since $TC_n(X) = secat(p)$ (see [8]), $cat(X^{n-1}) = secat(q)$ and $secat(q) \leq secat(p)$, we have

$$TC_n(X_0) \leq r(n-1) = cat(x_0^{n-1}) \leq TC_n(X_0).$$

□

Acknowledgements. I would like to thank Y. Félix and A. Murillo for without them this paper would have never been written.

References

[1] O. Cornea, G. Lupton, J. Oprea, D. Tanré, *Lusternik-Schnirelmann category.* Math. Surveys Monogr. **103**, AMS, 2003.
[2] M. Farber, Topological complexity of motion planning. Discrete Comput. Geom. **29** (2003), 211–221.
[3] A. Fassò Velenik, *Relative homotopy invariants of the type of the Lusternik-Schnirelmann category.* Eingereichte Dissertation (Ph.D. Thesis), Freien Universität Berlin, 2002.
[4] Y. Félix, S. Halperin and J.-C. Thomas, *Rational Homotopy Theory. Grad. Texts in Math.* **205**, Springer, 2000.
[5] Y. Félix, D. Tanré, Formalité d'une application et suite spectrale d'Eilemberg-Moore. Lect. Notes in Math. **1318** (1988), 99–123.
[6] B. Jessup, A. Murillo, P.-E. Parent, Rational Topological Complexity. Accepted in Algebr. Geom. Topol. 2012.
[7] L. Lechuga, A. Murillo, Topological complexity of formal spaces. Contemp. Math. **438** (2007), 105–114.
[8] Y.B. Rudyak, On higher analogs of topological complexity. Topology Appl. **157** (2010), 916–920.
[9] A.S. Schwarz, The genus of a fiber space. Amer. Math. Soc. Transl. **55** (1966), 49-140.
[10] D. Sullivan, Infinitesimal computations in Topology. *Publ. Math. de l'I.H.E.S.* **47** (1978), 269–331.
[11] M. Vigué-Poirrier, *Formalité d'une application continue.* C. R. Acad. Sci. Paris Sr. A-B. **289** (1979), no. 16, A809-A812

Institut de Recherche en Mathématique et Physique, Chemin du Cyclotron, 2, B-1348 Louvain-la-Neuve,, Belgium.,
E-mail address: jose.carrasquel@uclouvain.be